



ORIGINAL ARTICLE

Quality of vision in endoscopy in the midst of a pandemic: Does PPE influence quality of vision during gastrointestinal endoscopy?☆



Reinaldo Andrés Rincón Sánchez^{a,*}, Alejandro Concha Mejía^b, Liana Marcela Viaña Ríos^c

^a Pontificia Universidad Javeriana. Hospital Universitario San Ignacio. Universidad de la Sabana. Fundación Clínica Shaio, Bogotá, Colombia

^b Universidad de la Sabana. Fundación Clínica Shaio. Fundación Universitaria Sanitas. Clínica Colombia, Bogotá, Colombia

^c Pontificia Universidad Javeriana. Hospital Universitario San Ignacio. Organización Sura, Bogotá, Colombia

Received 24 July 2020; accepted 18 October 2020

Available online 14 September 2021

KEYWORDS

COVID-19;
Optical quality;
Personal protective
equipment;
Endoscopy

Abstract

Introduction: In the midst of the SARS-CoV2 virus (COVID-19) pandemic, health professionals, specifically gastroenterologists, have had to use personal protective equipment (PPE) to reduce contact with droplets and aerosols generated during gastrointestinal endoscopy.

Objective: To evaluate the impact of the use of two types of PPE on quality of vision during gastrointestinal endoscopy.

Methods: A cross-sectional observational pilot study in gastroenterologists who undergo an ophthalmological examination of visual acuity and quality of vision when using two types of PPE. Type #1: 3M N95 1860 green respirator + 3M mono safety glasses + protective screen. Type #2: 3M 6800 full facepiece + 3M NIOSH 7093C HF/P100 filters.

Results: Visual acuity and quality of vision parameters while using the PPE that is routinely used when performing gastrointestinal endoscopy during the pandemic were evaluated. It was found that Modality #1 was associated with decreases of up to 37% in visual acuity, 25% in colour visualisation and 75% in contrast sensitivity among digestive endoscopists within minutes of placement. These figures worsened over the course of the procedure, rising to 75%, 60% and 100%, respectively. Modality #2 was not associated with any deterioration in quality of vision.

Conclusions: The different PPE modalities used during gastrointestinal endoscopy could have an impact on the quality of endoscopy studies performed during the SARS COV2 (COVID-19) pandemic.

© 2020 Elsevier España, S.L.U. All rights reserved.

☆ Please cite this article as: Rincón Sánchez RA, Concha Mejía A, Viaña Ríos LM. Calidad de la visión en endoscopia en tiempos de pandemia: ¿Influyen los EPP en la calidad visual durante la endoscopia digestiva? Gastroenterología y Hepatología. 2021;44:637–643.

* Corresponding author.

E-mail address: rincon_r@javeriana.edu.co (R.A. Rincón Sánchez).

PALABRAS CLAVE

Covid-19;
Calidad óptica;
Elementos de
protección personal;
Endoscopia

Calidad de la visión en endoscopia en tiempos de pandemia: ¿Influyen los EPP en la calidad visual durante la endoscopia digestiva?

Resumen

Introducción: En tiempos de la pandemia por el virus SARS-CoV2 (Covid-19), los profesionales de la salud y específicamente los gastroenterólogos hemos tenido la necesidad de usar elementos de protección personal (EPP) para disminuir el contacto con microgotas y aerosoles durante la realización de endoscopia digestiva.

Objetivo: Evaluar el impacto del uso de dos modalidades de elementos de protección personal en la calidad de la visión durante la endoscopia digestiva.

Métodos: Estudio piloto observacional de corte transversal en gastroenterólogos a quienes se les realiza examen oftalmológico de agudeza y calidad visual al utilizar dos modalidades de EPP. Modalidad #1: Respirador N95 3M 1860 VERDE + Monogafas 3M + Pantalla protectora y modalidad #2: Pieza Facial de cara completa 3M 6800 con filtros 3M 7093C NIOSH HF/P100.

Resultados: Al evaluar los parámetros de agudeza y calidad visual al usar los EPP que se utilizan rutinariamente en el ejercicio de la endoscopia digestiva durante la pandemia, se identifica que con el uso de la modalidad #1, una disminución de hasta el 37% en la agudeza y en 25% en la visualización del color y el 75% en la sensibilidad al contraste en endoscopistas digestivos desde los primeros minutos de la colocación, lo cual empeora en el tiempo, hasta 75, 60 y 100% respectivamente durante el procedimiento. En el caso de la modalidad #2, no hay deterioro de la calidad de la visión.

Conclusiones: Las diferentes modalidades de EPP utilizados durante la endoscopia digestiva podrían tener repercusión en la calidad de los estudios endoscópicos realizados durante la pandemia por SARS COV2 (Covid-19).

© 2020 Elsevier España, S.L.U. Todos los derechos reservados.

Introduction and rationale

Since the advent of the SARS-CoV-2 (COVID-19) pandemic in December 2019, the entire population, particularly health personnel, has been forced to use personal protective equipment (PPE) to reduce the probability of contact and therefore of contagion of the disease caused by this agent, whose virulence is high compared to other recent epidemics in the world.¹

Specifically in digestive endoscopy, the production of droplets and the protection of the personnel directly involved in the procedures has acquired great relevance, leading to the discussion of the subject in numerous forums, both in gastroenterology and endoscopy associations and in health institutions' infectious disease committees. The recommendation made by most of the documents produced by such meetings and consensus is the use of respirators with N95 or higher filters (FFP2 and FFP3) and eye protection, by means of mono glasses and/or protective shields to reduce exposure of the conjunctiva or the mucosa to the virus. There are also fullface pieces with N95 filters and simultaneous face and eye protection on the market which have become an alternative for some specialist physicians.^{2,3}

All endoscopists are well acquainted with the discomfort, the limitation upon free air flow during breathing and the effects of these devices on facial skin. However, another

relevant aspect is the reduction in visual acuity and quality produced by misting in some of the adjustable mono glasses or the additional glare that could limit the effectiveness of the procedure at any moment, either due to the reduced capacity to detect and classify small lesions, or because certain therapeutic procedures, the efficacy and safety of which in many cases hinge on precision, are hampered or delayed.^{4,5}

For this reason, a quantification and comparison of visual parameters such as visual acuity (VA) is proposed, where VA is defined as the human visual system's capacity to make out, recognise or distinguish details in objects under conditions of high contrast and with a good level of illumination,⁶ in the Snellen fraction, contrast sensitivity (CS), which represents the lowest contrast the visual system can detect⁷ and the Ishihara colour vision test (Cv), a test that uses random-sized and coloured dot circles to identify colour vision defects, in endoscopist specialists with different PPE modalities, for the purpose of evaluating the impact of the use of two modalities of personal protective equipment on vision quality during digestive endoscopy.

Methods

An observational cross-sectional pilot study was carried out, in which gastroenterology professors with more than 10

Table 1 Data from the assessment of visual acuity, contrast and colour visualisation with the use of two modalities of personal protective equipment.

Subject 1. VA with corrected distance (OD: -4.25 SPH OS: -3.50 – 0.75 × 160) OD: 20/20 OS:20/20. CS: AO: 6%. CV: 10/10									Subject 2. VA without corrected distance OD: 20/20 OS:20/20 CS AO:6% CV: 10/10								
Time of use of PPE	PPE1			PPE2			DIFFERENCE ACCORD-ING TO MODAL-ITY %		Time of use of PPE	PPE1			PPE2			DIFFERENCE ACCORD-ING TO MODAL-ITY %	
	SNELLEN VA, CS, CV	DECIMAL SCALE		SNELLEN VA, CS, CV	DECIMAL SCALE		OD	OS		SNELLEN VA, CS, CV	DECIMAL SCALE		SNELLEN VA, CS, CV	DECIMAL SCALE		OD	OS
		OD	OS		OD	OS					OD	OS		OD	OS		
Minute 0	VA OD:20/30 OS: 20/50- CS: OD: -OS: -	0.63	0.40	VA OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	37%	60%	Minute 0	VA OD: 20/20 OS: 20/20 CS: OD: 25% OS: 25% CV: OD: 8/10 OS: 8/10	1	1.00	VA OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	0%	0%
		0	0		1	1	100%	100%			0.50	0.50		1	1	50%	50%
		0.70	0.30		1	1	30%	70%			0.80	0.80		1	1	20%	20%
Minute 15	VA OD: 20/80 OS: 20/100- CS: OD: -OS: -	0.25	0.20	VA OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	75%	80%	Minute 15	VA OD:20/200 OS: 20/80- CS: OD: -OS: -	0.10	0.25	VA OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	90%	75%
		0	0		1	1	100%	100%			0	0.00		1	1	100%	100%
		0.50	0.10		1	1	50%	90%			0.80	0.80		1	1	20%	20%
Minute 25	VA OD: 20/150 OS: 20/200- CS: OD: -OS: -	0.13	0.10	VA: OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	87%	90%	Minute 25	VA OD: 20/80 OS: 20/50- CS: OD: -OS: -	0.25	0.40	VA: OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	75%	60%
		0	0		1	1	100%	100%			0	0.00		1	1	100%	100%
		0.5	0.50		1	1	50%	100%			0.50	0.80		1	1	50%	20%
Minute 45	VA OD: 20/200 OS:20/200- CS: OD: -OS: -	0.10	0.10	VA: OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	90%	90%	Minute 45	VA OD: 20/150 OS: 20/80- CS: OD: -OS: -	0.13	0.25	VA: OD: 20/20 OS: 20/20 CS: OD: 6% OS: 6%	1	1	87%	75%
		0	0		1	1	100%	100%			0	0.00		1	1	100%	100%
		0	0		1	1	70%	100%			0.30	0.80		1	1	70%	20%

VA: visual acuity; CS: contrast sensitivity; CV: colour vision; OD: right eye; OS: left eye.



Figure 5 Assessment of subjects 1 and 2 with PPE 2.

an independent evaluation of both eyes of each previous subject, minute 0, 15, 25 and 45 (Table 1).

There is a significant decrease that worsens over time in the three parameters with the use of the first PPE modality (N95 3M 1860 green respirator + 3M mono glasses + protective shield), with a 37% reduction in visual acuity when the equipment is donned and of up to 75% after 45 minutes of use (Fig. 6), contrast sensitivity compromise is greater than 75% initially and is 100% after 45 minutes (Fig. 7), whereas in colour vision it is 25% at the beginning and up to 60% after 45 minutes (Fig. 8).

With the use of the second PPE modality (3M 6800 full facepiece with NIOSH 3M 7093C HF/P100 filters). There is no variation in any of the aforementioned parameters and visual quality remains equal to the baseline.

Discussion

This is the first study to evaluate the impact of different PPE modalities on visualisation quality in medical procedures during the SARS-CoV-2 (COVID-19) pandemic and which impact the assessment of optical parameters and probably the quality of endoscopic studies (defined as an increased rate of complications and a reduction in diagnostic performance for lesions with subtle changes in the mucosal pattern, whether they are preneoplastic or early neoplastic). This alteration is due to significant limitations in visual acuity, contrast sensitivity and colour visualisation. These parameters are affected in this study between the moment the PPE are donned and worsen over time depending on the duration of the procedure.

Full facepiece PPE do not alter the optical parameters evaluated in the first measurement or as the procedure evolves and are equal to the optical measurements obtained without the use of PPE, which is considered the best vision possible for each subject.

Studies with a greater number of measurements that could confirm or refute the findings of this pilot study are required, as are prospective studies that would make it

possible to determine whether the visual alterations documented during this pandemic have an impact on strong endoscopic quality outcomes.

One additional, but no less important, point, will be the evaluation of the impact on ergonomics and health at work PPE entailed in the use of this PPE.

Conclusion

Some PPE modalities used during gastrointestinal endoscopy reduce endoscopists' quality of vision, which could affect the quality and even the frequency of complications of diagnostic and therapeutic procedures performed during the SARS-CoV-2 (COVID-19) pandemic.

Funding

No funding was received for this study.

Conflicts of interest

The authors declare that they have no conflicts of interest.

References

1. World Health Organization. Rational use of personal protective equipment for coronavirus disease (COVID-19): interim guidance; 2020. Available from: <https://extranet.who.int/iris/restricted/handle/10665/33121539>.
2. Repici A, Maselli R, Colombo M, Gabbiadini R, Spadaccini M, Anderloni A, et al. Coronavirus (COVID19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc.* 2020;S0016510.
3. Otero W, Gomez M, Ángel LA, Ruiz O, Marulanda H, Riveros J, et al. Procedimientos endoscópicos y pandemia COVID-19. Consideraciones básicas. *Rev Col Gastroenterol.* 2020;35:65–75.
4. Rizk M, Sawhney M, Cohen J, Pike IM, Adler DG, Dominitz JA, et al. Quality Indicators Common to All GI Endoscopic Procedures. *Am J Gastroenterol.* 2015;110:48–59.




Figure 6




Figure 7




Figure 8

5. López-Picazo J. Quality indicators in digestive endoscopy: introduction to structure, process, and outcome common indicators. *Rev Esp Enferm Dig.* 2017;109: 435–50.
6. Montés-Micó R. *Optometría: Principios básicos y aplicación clínica.* Barcelona: Elsevier España; 2011. p. 105.
7. López Y. Importancia de la valoración de sensibilidad al contraste en la práctica optométrica. *CTSVO.* 2009;7:99–114.